




FULL - DOSE STANOZOLOL

Concentration ____ 10mg/Tablet

Route of A. _____ Oral

Schedule _____ CIII - Rx Only

Form _____ Tablet

Stanozolol is an orally administered anabolic-androgenic steroid derived from dihydrotestosterone (DHT). It is a 17-alpha-alkylated compound formulated to remain active after gastrointestinal absorption. Due to its pharmacological structure, stanozolol exhibits anabolic activity with relatively lower androgenic properties compared with some other androgenic agents, although the overall clinical response may vary according to dose, duration of use, and individual patient factors.

Stanozolol may influence protein metabolism and contribute to nitrogen retention. In medical use, it has been associated with selected therapeutic applications, including certain conditions involving tissue breakdown and, historically, hereditary angioedema. However, therapeutic response may differ depending on the patient's general health status, underlying disease profile, nutritional condition, and concomitant treatments.

As an orally active anabolic agent, stanozolol should be used under careful medical supervision because of its potential effects on hepatic function, lipid metabolism, endocrine balance, and cardiovascular parameters. Periodic clinical and laboratory monitoring, including liver function tests and other relevant evaluations, should be considered during treatment.

KEEP OUT OF REACH OF CHILDREN
STORE TABLET IN COOL,
DRY PLACE BETWEEN 15-25°C (59-77°F)

PROTECT TABLET FROM LIGHT AND
MOISTURE USE TABLET ONLY AS DIRECTED
BY A HEALTHCARE PROFESSIONAL

Norwick



ACCREDITED



ONLY

norwickpharma.com
hi@norwickpharma.com

88 The Parq Building, Ratchadaphisek Road
Klongtoey Sub-district, Bangkok / Thailand